

Editorial Comment

Progressive Inadequacy of Vascular Support in Myocardial Hypertrophy*

DAVID H. SPODICK, MD, DSc, FACC

Worcester, Massachusetts

Hypertrophy: advantages and eventual disadvantages.

Unlike other organs subject to hemodynamic overload, the myocardium responds by hypertrophy; survival compels coronary vascular adaptation. Left ventricular hypertrophy, the common structural response to a variety of overloads, can be compensatory early but detrimental when it progresses without effective treatment. Left ventricular hypertrophy initially decreases wall stress but ultimately induces contraction and relaxation abnormalities and vulnerability to arrhythmias, particularly if it is associated with coronary disease. Moreover, although left ventricular hypertrophy may limit infarct expansion (1), infarcts are larger in hypertrophic ventricles (2), perhaps because of inadequate growth factors for angiogenesis of collateral channels (3). Even with normal epicardial arteries, the vascular hierarchy responds in proportion to the severity and duration of hypertrophy and its pathogenetic process. Consequently, left ventricular hypertrophy that exceeds its vascular support at any level becomes ischemic; moreover, left ventricular hypertrophy disproportionate to diastolic volume permits wall stress to increase unchecked and, with it, oxygen demand, independent of vascular conditions.

Myocardial/microvascular flow disproportion. Myocardial and vascular growth need not proceed in parallel. Although myocardial fibers hypertrophy as a cube function and capillary surface increases only as a square function, early left ventricular hypertrophy may be adequate for demands during rest. Yet increased total coronary dimensions and flow, initially compensatory, are ultimately outpaced by increasing ventricular mass, reducing flow per unit of myocardium (4,5). In the microvasculature, capillary diameters may be similar in normal and hypertrophic ventricles, but *total capillary surface decreases in left ventricular hypertrophy* and, despite normal arteriolar density (6), greater intercapillary separation (7) increases distances for

blood-myocardial exchange of nutrients and metabolites. Extravascular compression by hypertrophic muscle and decreased capacity for maximal vasodilation exacerbate these disadvantages. The resultant *decreased coronary reserve* is minimal in early left ventricular hypertrophy and is often evident only as a response to physiologic or paced tachycardia or to hypoxic or induced (e.g., adenosine) vasodilator stimuli, or as abnormal reactive hyperemia; it worsens disproportionately with progressive left ventricular hypertrophy. Decreased production of endothelium-derived relaxing factor may contribute (8) as may inappropriate vasoconstriction of prearteriolar vessels (9); both exacerbate the increased arteriolar wall to lumen ratio in left ventricular hypertrophy (10) with abnormal shear stress at the blood-endothelial interface (11). Even in "pure" left ventricular hypertrophy, increased left ventricular mass alone greatly increases coronary resistance and decreases flow reserve (maximal/rest flow ratio). Coronary venous compression probably adds to total coronary resistance.

Any increased cavity pressure will also compress sub-endocardial vessels, whereas abnormal myocardial relaxation additionally tends to compress intramural coronary vessels, further shifting transmural blood flow (12,13). Because 80% of left ventricular coronary flow is diastolic, prolonged systole, due mainly to prolonged left ventricular ejection time in compensated aortic stenosis (14), prolongs "physiologic" systolic ischemia and encroaches on diastolic time for flow and metabolic recuperation (for example, time for cytosolic calcium ion to return to the sarcoplasmic reticulum). Thus, the normally relatively ischemic subendocardial circulation is further hypoperfused in left ventricular hypertrophy during exercise, tachycardia and vasodilator stimulation, with a more unfavorable shift toward the mid- and subepicardial layers (15) contributing to any clinical ischemia.

Vasculature in aortic stenosis. In aortic stenosis normal coronary conduit arteries may permit appropriate left ventricular hypertrophy with normal cavity size and better function (16). Yet minor increases in epicardial coronary resistance affect total coronary resistance very little compared with the great *selective decrease of microvascular reserve* in severe aortic stenosis with left ventricular hypertrophy. The latter, a mechanism for angina with unobstructed epicardial coronary arteries, can be conceived as a special case of "syndrome X." Thus, major coronary narrowing could be disastrous because low poststenotic coronary pressure exacerbates the downstream flow-limiting effects of extravascular compression (17) and the other determinants of decreased coronary reserve, whereas enlargement of conduit arteries, even proportional to myocardial hypertrophy, would have little effect on microvascular ischemia.

Vasculature in volume and pressure overloading: present study. Volume and pressure loading by either aortic regurgitation or stenosis yield grossly similar directional effects on

*Editorials published in *Journal of the American College of Cardiology* reflect the views of the authors and do not necessarily represent the views of JACC or the American College of Cardiology.

From the Cardiology Division, St. Vincent Hospital and the Department of Medicine, University of Massachusetts Medical School, Worcester, Massachusetts.

Address for correspondence: David H. Spodick, MD, Cardiology Division, St. Vincent Hospital, Worcester, Massachusetts 01604.

coronary reserve (18); systemic hypertension also does so (19) despite inclusion of the coronary orifices that are outside the overload zone of aortic valve disease. Intuitively, however, left ventricular hypertrophy differing in pathogenesis should have different associated effects at some levels. For example, patients with normal epicardial coronary arteries have much more angina with aortic stenosis than with aortic or mitral regurgitation and their respective coronary reserves differ correspondingly (20). Because the reduction in coronary reserve associated with left ventricular hypertrophy tends to be greater in humans than in experimental animals (21), the report by Villari et al. (22) in this issue of the Journal has dual importance as it involves both human subjects and progressive levels of left ventricular hypertrophy. The authors (22) might have reported on patients with aortic stenosis and regurgitation differentially, but they did not investigate the microvasculature. Nevertheless, their subjects, all with unobstructed epicardial coronary arteries, were selected because, with the exception of increased left anterior descending and left circumflex artery cross-sectional area, they were comparable at baseline to control subjects who underwent cardiac catheterization for chest pain. This increased cross-sectional area, when normalized for left ventricular mass, was initially comparable to that of control subjects, but it decreased significantly, becoming disproportionate as the patients' condition progressed from initially moderate to subsequent marked left ventricular hypertrophy.

In conclusion, progressively inadequate myocardial microvascular support converts progressive left ventricular hypertrophy from a transiently beneficial adaptation to an ultimately detrimental process. That the macrovascular circulation also fails to keep pace with myocardial growth requires further investigation.

References

1. Pirolo JS, Hutchins GM, Moore GW. Infarct expansion: pathologic analysis of 204 patients with a single myocardial infarct. *J Am Coll Cardiol* 1986;7:249-54.
2. Koyanagi S, Eastham CL, Harrison DG, Marcus ML. Increased size of myocardial infarction in dogs with chronic hypertension and LVH. *Circ Res* 1982;50:55-62.
3. Sasayama S, Fujita M. Recent insights into coronary collateral circulation. *Circulation* 1992;85:1197-204.
4. Roberts CS, Roberts WC. Cross-sectional area of the proximal portions of the three major epicardial coronary arteries in 98 necropsy patients with different coronary events: relationship to heart weight, age and sex. *Circulation* 1980;62:953-9.
5. Schwitzer J, Eberli FR, Ritter M, Turina M, Krayenbuehl HP. Myocardial oxygen consumption in aortic valve disease with and without LV dysfunction. *Br Heart J* 1992;67:161-9.
6. Tomanek RJ, Palmer PJ, Pfeiffer GL, Schreiber KL, Eastham CL, Marcus ML. Morphometry of canine coronary arteries, arterioles and capillaries during hypertension and left ventricular hypertrophy. *Circ Res* 1986;58:38-46.
7. Breisch EA, Houser SR, Carey RA. Myocardial blood flow and capillary density in chronic pressure overload of the feline LV. *Cardiovasc Res* 1980;14:469-75.
8. Maseri A, Crea F, Kaski JC, Crake T. Mechanisms of angina pectoris in syndrome X. *J Am Coll Cardiol* 1991;17:499-506.
9. Bassenge E, Busse R. Endothelial modulation of vascular tone. *Prog Cardiovasc Dis* 1988;30:349-80.
10. Marcus ML, Mueller TM, Eastham CL. Effects of short and long term left ventricular hypertrophy on coronary circulation. *Am J Physiol* 1981;241:H358-62.
11. Stack RS, Rembert JC, Schirmer B, Greenfield JC Jr. Relation of left ventricular mass to geometry of the proximal coronary arteries in the dog. *Am J Cardiol* 1983;51:1728-31.
12. Murakami T, Hess OM, Gage JE, Grimm J, Krayenbuehl HP. Diastolic filling dynamics in patients with aortic valve disease. *Circulation* 1986;73:1162-74.
13. Alyono D, Anderson RW, Parrish DG, Dai X-Z, Bache RJ. Alterations of myocardial blood flow associated with experimental canine LVH secondary to aortic stenosis. *Circ Res* 1986;58:47-57.
14. Spodick DH. Systolic time intervals reconsidered. *Am J Cardiol* 1984;53:1667-70.
15. Vatner SF, Shannon R, Hittinger L. Reduced subendocardial coronary reserve. A potential mechanism for impaired diastolic function in the hypertrophied and failing heart. *Circulation* 1990;81(suppl II):III-8-14.
16. Vekshtein V, Alexander RW, Plappert T, et al. Coronary artery disease is associated with inadequate left ventricular hypertrophy in aortic stenosis (abstr). *J Am Coll Cardiol* 1989;14(suppl A):81A.
17. Bache RJ, Wright L, Laxson DD, Dai X-Z. Effect of coronary stenosis on myocardial blood flow during exercise in the chronically pressure-overloaded hypertrophied left ventricle. *Circulation* 1990;81:1967-73.
18. Marcus ML, Mueller TM, Gascho JA, Kerber RE. Effects of cardiac hypertrophy secondary to hypertension on the coronary circulation. *Am J Cardiol* 1979;44:1023-8.
19. Tubau JF, Szlachet J, London MJ, Hollenberg M, Mangano D, Massie BM. Systemic hypertension, LVH and coronary artery disease. *Am J Cardiol* 1979;44:1023-8.
20. Hiratzka LF, Doty BD, Eastham CL, Marcus ML, White CW. Pressure versus volume overload has different effects on coronary reserve (abstr). *Circulation* 1982;66(suppl II):II-354.
21. Marcus ML, Harrison DG, Chilian WM, et al. Alterations in the coronary circulation in hypertrophied ventricles. *Circulation* 1987;75(suppl I):I-19-25.
22. Villari B, Hess OM, Moccetti D, Vassalli G, Krayenbuehl HP. Effect of progression of left ventricular hypertrophy on coronary artery dimensions in aortic valve disease. *J Am Coll Cardiol* 1992;20:1073-9.